

# Revised Iterative Solution of Ground State of Double-Well Potential

Zhao Wei-Qin<sup>1, 2</sup>

1. *China Center of Advanced Science and Technology (CCAST)*  
*(World Lab.), P.O. Box 8730, Beijing 100080, China*
2. *Institute of High Energy Physics, Chinese Academy of Sciences,*  
*P. O. Box 918(4-1), Beijing 100039, China*

## Abstract

A revised new iterative method based on Green function defined by quadratures along a single trajectory is developed and applied to solve the ground state of the double-well potential. The result is compared to the one based on the original iterative method. The limitation of the asymptotic expansion is also discussed.

PACS: 11.10.Ef, 03.65.Ge

Key words: iterative solution, asymptotic expansion, double-well potential

# 1. Introduction

The double-well potential in one dimension,

$$V(x) = \frac{1}{2}g^2(x^2 - 1)^2, \quad (1.1)$$

is specially interesting since it has degenerate minima and could be served as a simple example of the bound-state tunnelling problem in quantum mechanics. However, it is a non-perturbative problem to solve the Schroedinger equation for this potential due to the tunnelling effect between the two minima. The asymptotic series of the ground state energy is meaningful only for quite large  $g$  (e.g.  $g \geq 6$ ). Even for such large  $g$  the obtained plateau in the energy expansion series does not necessarily consistent to the exact solution. An effective method to obtain the convergent solution of this problem for any values of  $g$  is needed.

Recently an iterative solution of the ground state for the one dimensional double-well potential is obtained[1] based on the Green function method developed in ref.[2]. This Green function is defined along a single trajectory, from which the ground state wave function in N-dimension can be expressed by quadratures along the single trajectory. This makes it possible to develop an iterative method to obtain the ground state wave function, starting from a properly chosen trial function. The convergence of the iterative solution very much depends on the choice of the trial function[1].

However, in the original iterative solution the obtained correction for the trial wave function is in the form of a power expansion, while the bound state solution should be in the form of an exponential when the coordinate variable approaches infinity. Recently a new revised iterative procedure[3] based on the same Green function is developed and applied to solve anhamornic oscillator and Stark effect. This method has some advantages compared to the original one. It not only gives an exponential form for the correction of the trial wave function, but also spends much less time in each iteration due to less folds of integration. It is natural to try this method to solve the double-well potential which has been attracted much attention.

In Section 2, a brief introduction is given about the Green function method based on the single trajectory quadrature. Special discussion is given to the revision of the iterative formula. The revised iterative formula for the double-well potential is given in Section 3, together with the trial function and the boundary condition for the lowest even state. The numerical results based on the revised iterative formula and the original iterative method are collected in Section 4, together with a comparison to the asymptotic expansion. A compact expression of the asymptotic expansion is derived in Appendix A. Finally some discussions are given at the end.

## 2. Green Function and the Revised Iterative Solution

For a particle with unit mass, moving in an N-dimensional unperturbed potential  $V_0(\mathbf{q})$ , the ground state wave function  $\Phi(\mathbf{q})$  satisfies the following Schroedinger equation:

$$H\Phi(\mathbf{q}) = E\Phi(\mathbf{q}), \quad (2.1)$$

where

$$H = T + V_0(\mathbf{q}) = -\frac{1}{2}\nabla^2 + V_0(\mathbf{q}). \quad (2.2)$$

Assume the solution of eq.(2.1) could be expressed as

$$\Phi(\mathbf{q}) = e^{-S(\mathbf{q})}. \quad (2.3)$$

Introduce a perturbed potential  $U(\mathbf{q})$  and assume

$$V(\mathbf{q}) = V_0(\mathbf{q}) + U(\mathbf{q}) \quad (2.4)$$

$$\mathcal{H} = T + V(\mathbf{q}) = -\frac{1}{2}\nabla^2 + V(\mathbf{q}). \quad (2.5)$$

Define another wave function  $\Psi(\mathbf{q})$  satisfying the Schroedinger equation

$$\mathcal{H}\Psi(\mathbf{q}) = \mathcal{E}\Psi(\mathbf{q}) \quad (2.6)$$

$$\mathcal{E} = E + \Delta. \quad (2.7)$$

Let

$$\Psi(\mathbf{q}) = e^{-S(\mathbf{q})-\tau(\mathbf{q})}. \quad (2.8)$$

The equation for  $\tau$  and  $\Delta$  could be derived easily[2]:

$$\nabla S \cdot \nabla \tau + \frac{1}{2}[(\nabla \tau)^2 - \nabla^2 \tau] = (U - \Delta). \quad (2.9)$$

Consider the coordinate transformation

$$(q_1, q_2, q_3, \dots, q_N) \rightarrow (S, \alpha_1, \alpha_2, \dots, \alpha_{N-1}) = (S, \alpha) \quad (2.10)$$

with  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{N-1})$  denoting the set of  $N - 1$  orthogonal angular coordinates satisfying the condition

$$\nabla S \cdot \nabla \alpha_i = 0, \quad (2.11)$$

for  $i = 1, 2, \dots, N-1$ . Some useful quantities for this coordinate transformation[2] are listed in Appendix B. Similarly to the discussions in Ref.[2], introducing the  $\theta$ -function in S-space:

$$\theta(S - \bar{S}) = \begin{cases} 1 & \text{if } 0 \leq \bar{S} < S \\ 0 & \text{if } 0 \leq S < \bar{S} \end{cases} \quad (2.12)$$

and define

$$C = \theta[(\nabla S)^2]^{-1} = \theta h_S^2. \quad (2.13)$$

Using

$$\nabla S \cdot \nabla C = 1, \quad (2.14)$$

it is easy to derive the following equation

$$\tau = (1 + CT)^{-1} C [(U - \Delta) - \frac{1}{2}(\nabla \tau)^2]. \quad (2.15)$$

When the N-dimensional variable  $\mathbf{q}$  is transformed into  $(S, \alpha)$ ,  $T = -\frac{1}{2}\nabla^2$  could be decomposed into two parts:

$$T = T_S + T_\alpha, \quad (2.16)$$

where  $T_S$  and  $T_\alpha$  consist only the differentiation to  $S$  and  $\alpha$ , respectively. The detailed expression of  $T_S$  and  $T_\alpha$  is given in Appendix B. Now another Green function could be defined as[2]

$$\bar{D} \equiv -2\theta e^{2S} \frac{h_S}{h_\alpha} \theta e^{-2S} h_S h_\alpha \quad (2.17)$$

and it is related to  $C$  in the following way[2]:

$$(1 + \bar{D}T_\alpha)^{-1} \bar{D} = (1 + CT)^{-1} C. \quad (2.18)$$

Therefore, from (2.15), we have

$$\tau = (1 + \bar{D}T_\alpha)^{-1} \bar{D} [(U - \Delta) - \frac{1}{2}(\nabla \tau)^2]. \quad (2.19)$$

The explicit expression of  $\tau$  based on (2.19) is

$$\tau = -2 \int_0^S e^{2S'} \frac{h_{S'}}{h_\alpha} dS' \int_0^{S'} e^{-2S''} h_{S''} h_\alpha dS'' (1 + T_\alpha \bar{D})^{-1} [(U - \Delta) - \frac{1}{2}(\nabla \tau)^2]. \quad (2.20)$$

Therefore, we have

$$-\frac{1}{2} \frac{h_\alpha}{h_S} e^{-2S} \frac{\partial \tau(S, \alpha)}{\partial S} = \int_0^S e^{-2S'} h_{S'} h_\alpha dS' (1 + T_\alpha \bar{D})^{-1} [(U - \Delta) - \frac{1}{2}(\nabla \tau)^2]. \quad (2.21)$$

The left hand side of Eq.(2.20) approaches to 0 when  $S \rightarrow \infty$ , so is the right hand side, i.e.,

$$\int_0^\infty e^{-2S} h_S h_\alpha dS (1 + T_\alpha \overline{D})^{-1} [(U - \Delta) - \frac{1}{2}(\nabla\tau)^2] = 0, \quad (2.22)$$

which is correct for all  $\alpha$ . Integrating over  $d\alpha = \Pi_{i=1}^{N-1} d\alpha_i$  and because of

$$h_S h_\alpha T_\alpha = -\frac{1}{2} \sum_{j=1}^{N-1} \frac{\partial}{\partial \alpha_j} \frac{h_S h_\alpha}{h_j^2} \frac{\partial}{\partial \alpha_j} \quad \text{and} \quad \int d\alpha h_S h_\alpha T_\alpha \tau = 0, \quad (2.23)$$

we derive

$$\int h_S h_\alpha d\alpha dS e^{-2S} [(U - \Delta) - \frac{1}{2}(\nabla\tau)^2] = 0. \quad (2.24)$$

Denoting  $d\mathbf{q} = h_S h_\alpha d\alpha dS$ , from Eq.(2.3), we reach a new expression of the perturbative energy

$$\Delta = \frac{\int d\mathbf{q} \Phi^2 [U - \frac{1}{2}(\nabla\tau)^2]}{\int d\mathbf{q} \Phi^2}. \quad (2.25)$$

Based on Eqs. (2.25) and (2.15) or (2.19) we have the new iteration series

$$\begin{aligned} \Delta_n &= \frac{\int d\mathbf{q} \Phi^2 [U - \frac{1}{2}(\nabla\tau_{n-1})^2]}{\int d\mathbf{q} \Phi^2}, \\ \tau_n &= (1 + \overline{D}T_\alpha)^{-1} \overline{D} [(U - \Delta_n) - \frac{1}{2}(\nabla\tau_{n-1})^2] \\ \text{or} \\ \tau_n &= (1 + CT)^{-1} C [(U - \Delta_n) - \frac{1}{2}(\nabla\tau_{n-1})^2]. \end{aligned} \quad (2.26)$$

For later convenience this new iteration series is name as the  $\tau$ -iteration in this paper. Let us compare the above  $\tau$ -iteration with the original one derived from the equation for  $f = e^{-\tau}$  and  $\Delta$  in Ref.[2], which is named as  $f$ -iteration in this paper:

$$\begin{aligned} \Delta_n &= \frac{\int d\mathbf{q} \Phi^2 U f_{n-1}}{\int d\mathbf{q} \Phi^2 f_{n-1}} \\ f_n &= 1 + (1 + \overline{D}T_\alpha)^{-1} \overline{D} (-U + \Delta_n) f_{n-1} \\ \text{or} \\ f_n &= 1 + (1 + CT)^{-1} C (-U + \Delta_n) f_{n-1}. \end{aligned} \quad (2.27)$$

There are several advantages for the  $\tau$ -iteration:

- 1) It directly gives an exponential form for the perturbed wave function  $e^{-\tau}$ . This result is consistent with those obtained using the series expansion of  $\{\mathbf{S}_i\}$  and  $\{E_i\}$  (See Section 1 of Ref.[1]);
- 2) The iteration process in this formula is more transparent;
- 3) The calculation of the perturbation energy is much simpler.

### 3. Revised Iterative Formula for the Double-well Potential

In this section the revised iterative formula is applied to solve the ground state for the double-well potential. For the Hamiltonian  $H = T + V$  let us introduce the wave function for the lowest even eigenstate as  $\psi_{ev}$ , satisfying

$$H\psi_{ev} = E_{ev}\psi_{ev}. \quad (3.1)$$

In the following we are going to introduce the trial wave function  $\phi_{ev}$  for this state, satisfying

$$(H + w_{ev})\phi_{ev} = (E_{ev} + \mathcal{E}_{ev})\phi_{ev} = g\phi_{ev} \quad (3.2)$$

Assuming

$$\psi_{ev} = \phi_{ev}e^{-\tau_{ev}}, \quad (3.3)$$

the final energy and wave function  $\{\psi_{ev}, E_{ev}\}$  could be obtained by solving the corresponding equation of  $\{\tau_{ev}, \mathcal{E}_{ev}\}$  based on the revised iteration method introduced in Section 2. The key to choose the proper trial function is to satisfy the necessary boundary conditions of the state. For the ground state, namely the lowest even state, we have

$$\begin{aligned} \psi_{ev}(-x) &= \psi_{ev}(x) \\ \psi'_{ev}(0) &= 0 \\ \psi_{ev}(\infty) &= 0. \end{aligned} \quad (3.4)$$

The trial wave function should satisfy similar boundary conditions, namely,

$$\begin{aligned} \phi_{ev}(-x) &= \phi_{ev}(x) \\ \phi'_{ev}(0) &= 0 \\ \phi_{ev}(\infty) &= 0. \end{aligned} \quad (3.5)$$

Following the steps in Section 2 of Ref.[1], we introduce, for  $x \geq 0$ ,

$$\begin{aligned}
gS_0(x) &\equiv \frac{g}{3}(x-1)^2(x+2), \\
S_1(x) &\equiv \ln \frac{x+1}{2}, \\
\phi_+(x) = \phi_+(-x) &\equiv e^{-gS_0(x)-S_1(x)} = e^{-gS_0(x)}\left(\frac{2}{1+x}\right), \\
\phi_-(x) &\equiv e^{-gS_0(-x)-S_1(x)} = e^{-\frac{4}{3}g}e^{+gS_0(x)}\left(\frac{2}{1+x}\right).
\end{aligned} \tag{3.6}$$

It is easy to see that  $\phi_+(x)$  satisfies

$$(T + V + u)\phi_+ = g\phi_+, \tag{3.7}$$

where

$$\begin{aligned}
V(x) &= \frac{1}{2}g^2(x^2-1)^2 \\
u(x) &= \frac{1}{(1+x)^2}.
\end{aligned} \tag{3.8}$$

Following the necessary boundary conditions (3.5), we could choose the trial function as

$$\phi_{ev}(x) = \phi_{ev}(-x) \equiv \begin{cases} \phi_+(x) + \frac{g-1}{g+1}\phi_-(x), & \text{for } 0 \leq x < 1 \\ (1 + \frac{g-1}{g+1}e^{-\frac{4}{3}g})\phi_+(x), & \text{for } x > 1. \end{cases} \tag{3.9}$$

It is easy to proof that the above trial function satisfies Eq.(3.2) with

$$\begin{aligned}
w_{ev}(x) &= w_{ev}(-x) = u(x) + \hat{g}_{ev}(x), & \text{for } x \geq 0 \\
\hat{g}_{ev}(x) &= \begin{cases} 2g \frac{(g-1)e^{2gS_0(x)-\frac{4}{3}g}}{(g+1)+(g-1)e^{2gS_0(x)-\frac{4}{3}g}}, & \text{for } 0 \leq x < 1 \\ 0 & \text{for } x > 1. \end{cases}
\end{aligned} \tag{3.10}$$

Although the potentials  $V + w_{ev}$  is not continuous at  $x = 1$ , it could be proved that  $\phi_{ev}$  and  $\phi'_{ev}$  are continuous at any  $x$ , including  $x = 1$  and  $x = 0$ .

Now introduce the  $\theta$ -function in  $x$ -space:

$$(x|\theta|y) = \begin{cases} 0 & \text{for } x > y, \\ -1 & \text{for } x < y \end{cases} \tag{3.11}$$

and define the Green function

$$\overline{D}_{ev} \equiv -2\theta\phi_{ev}^{-2}\theta\phi_{ev}^2. \tag{3.12}$$

Following similar steps as in Section 2, we obtain the equations for  $\{\tau_{ev}, \mathcal{E}_{ev}\}$  as follows:

$$\begin{aligned}\tau_{ev} &= \overline{D}_{ev}[-(w_{ev} - \mathcal{E}_{ev}) - \frac{1}{2}(\tau'_{ev})^2], \\ \mathcal{E}_{ev} &= \frac{\int_0^\infty \phi_{ev}^2 (w_{ev} + \frac{1}{2}(\tau'_{ev})^2) dx}{\int_0^\infty \phi_{ev}^2 dx}.\end{aligned}\tag{3.13}$$

The solution of the ground state is

$$\psi_{ev} = \phi_{ev} e^{-\tau_{ev}}, \quad E_{ev} = g - \mathcal{E}_{ev}.\tag{3.14}$$

Similar procedure could also be taken by introducing

$$\overline{D}_+ \equiv -2\theta\phi_+^{-2}\theta\phi_+^2\tag{3.15}$$

and

$$\begin{aligned}\tau_+ &= \overline{D}_+[-(u - \mathcal{E}_+) - \frac{1}{2}(\tau'_+)^2], \\ \mathcal{E}_+ &= \frac{\int_0^\infty \phi_+^2 (u + \frac{1}{2}(\tau'_+)^2) dx}{\int_0^\infty \phi_+^2 dx}.\end{aligned}\tag{3.16}$$

It should be noticed that although  $\psi_+ = \phi_+ e^{-\tau_+}$  does satisfy the equation

$$H\psi_+ = E_+\psi_+,\tag{3.17}$$

with  $E_+ = g - \mathcal{E}_+$ , the solution  $\psi_+$  is not the eigenstate of the Hamiltonian  $H = T + V$ , because the trial function  $\phi_+$ , as well as the solution  $\psi_+$  do not satisfy the necessary boundary conditions. However, we still like to keep this solution here since an analytic expression of  $E_+$  and  $\psi_+$  has been obtained in terms of the asymptotic power expansion of  $1/g$  in Ref.[1]. This makes it possible to check the accuracy of the iteration procedure. In Appendix A compact expressions of the asymptotic power expansion of  $E_+$  and  $\tau'_+$  are derived.

Based on the explicit integral expression of  $\overline{D}_{ev}$  and  $\overline{D}_+$ , taking derivatives of  $\tau_{ev}$  and  $\tau_+$  in the first equations of (3.13) and (3.16), we obtain

$$\begin{aligned}\tau'_{ev}(x) &= 2\phi_{ev}^{-2}(x) \int_0^x \phi_{ev}^2(y) [(w_{ev}(y) - \mathcal{E}_{ev}) + \frac{1}{2}(\tau'_{ev}(y))^2] dy, \\ \mathcal{E}_{ev} &= \frac{\int_0^\infty \phi_{ev}^2(x) (w_{ev}(x) + \frac{1}{2}(\tau'_{ev}(x))^2) dx}{\int_0^\infty \phi_{ev}^2(x) dx}.\end{aligned}\tag{3.18}$$



and

$$\begin{aligned}\tau'_+(x) &= 2\phi_{ev}^{-2}(x) \int_0^x \phi_{ev}^2(y) [(u(y) - \mathcal{E}_+) + \frac{1}{2}(\tau'_+(y))^2] dy, \\ \mathcal{E}_+ &= \frac{\int_0^\infty \phi_+^2(x) (u(x) + \frac{1}{2}(\tau'_+(x))^2) dx}{\int_0^\infty \phi_+^2(x) dx}.\end{aligned}\tag{3.19}$$

The two sets of equations (3.18) and (3.19) are for the pairs of  $\tau'_{ev}$ ,  $\mathcal{E}_{ev}$  and  $\tau_+$ ,  $\mathcal{E}_+$ , which could be solved iteratively in the following way: Introducing the initial conditions

$$\mathcal{E}_{ev,0} = 0, \quad \tau'_{ev,0} = 0\tag{3.20}$$

and

$$\mathcal{E}_{+,0} = 0, \quad \tau'_{+,0} = 0,\tag{3.21}$$

we have, for  $n = 1$ ,

$$\begin{aligned}\mathcal{E}_{ev,1} &= \frac{\int_0^\infty \phi_{ev}^2(x) w_{ev}(x) dx}{\int_0^\infty \phi_{ev}^2(x) dx} \\ \tau'_{ev,1}(x) &= 2\phi_{ev}^{-2}(x) \int_0^x \phi_{ev}^2(y) (w_{ev}(y) - \mathcal{E}_{ev,1}) dy \\ &= -2\phi_{ev}^{-2}(x) \int_x^\infty \phi_{ev}^2(y) (w_{ev}(y) - \mathcal{E}_{ev,1}) dy\end{aligned}\tag{3.22}$$

and

$$\begin{aligned}\mathcal{E}_{+,1} &= \frac{\int_0^\infty \phi_+^2(x) u(x) dx}{\int_0^\infty \phi_+^2(x) dx}. \\ \tau'_{+,1}(x) &= 2\phi_+^{-2}(x) \int_0^x \phi_+^2(y) (u(y) - \mathcal{E}_{+,1}) dy \\ &= -2\phi_+^{-2}(x) \int_x^\infty \phi_+^2(y) (u(y) - \mathcal{E}_{+,1}) dy\end{aligned}\tag{3.23}$$

For  $n > 1$  we have

$$\mathcal{E}_{ev,n} = \mathcal{E}_{ev,1} + \frac{\int_0^\infty \phi_{ev}^2(x) \frac{1}{2}(\tau'_{ev,n-1}(x))^2 dx}{\int_0^\infty \phi_{ev}^2(x) dx} \quad (3.24)$$

$$\begin{aligned} \tau'_{ev,n}(x) &= \tau'_{ev,1}(x) + 2\phi_{ev}^{-2}(x) \int_0^x \phi_{ev}^2(y) [(\mathcal{E}_{ev,1} - \mathcal{E}_{ev,n}) + \frac{1}{2}(\tau'_{ev,n-1}(y))^2] dy \\ &= \tau'_{ev,1}(x) - 2\phi_{ev}^{-2}(x) \int_x^\infty \phi_{ev}^2(y) [(\mathcal{E}_{ev,1} - \mathcal{E}_{ev,n}) + \frac{1}{2}(\tau'_{ev,n-1}(y))^2] dy \end{aligned}$$

and

$$\mathcal{E}_{+,n} = \mathcal{E}_{+,1} + \frac{\int_0^\infty \phi_+^2(x) \frac{1}{2}(\tau'_{+,n-1}(x))^2 dx}{\int_0^\infty \phi_+^2(x) dx}. \quad (3.25)$$

$$\begin{aligned} \tau'_{+,n}(x) &= \tau'_{+,1}(x) + 2\phi_+^{-2}(x) \int_0^x \phi_+^2(y) [(\mathcal{E}_{+,1} - \mathcal{E}_{+,n}) + \frac{1}{2}(\tau'_{+,n-1}(y))^2] dy \\ &= \tau'_{+,1}(x) - 2\phi_+^{-2}(x) \int_x^\infty \phi_+^2(y) [(\mathcal{E}_{+,1} - \mathcal{E}_{+,n}) + \frac{1}{2}(\tau'_{+,n-1}(y))^2] dy \end{aligned}$$

For comparison we list in the following also the  $f$ -iteration formula for the double-well potential based on (2.27)[2]:

$$\begin{aligned} f_n(x) &= 1 - 2 \int_x^\infty \phi^{-2}(y) dy \int_y^\infty \phi^2(z) (w(z) - \mathcal{E}_n) f_{n-1}(z) dz \\ \mathcal{E}_n &= \int_0^\infty \phi^2(x) f_{n-1}(x) w(x) dx \Big/ \int_0^\infty \phi^2(x) f_{n-1}(x) dx, \end{aligned} \quad (3.26)$$

where  $f_n$  and  $\mathcal{E}_n$  could be either for the even ground state "ev" or the "+" state.

In the following section we are going to show the iterative results of  $\mathcal{E}_{+,n}$  and  $\mathcal{E}_{ev,n}$  based on the  $\tau$ -iteration formula Eqs.(3.24) and (3.25), together with the result from the  $f$ -iterative formula (3.26). The obtained  $\mathcal{E}_{+,n}$  is also compared to the asymptotic expansion for different  $g$ , obtained from the compact expression in Appendix A.

## 4. Numerical Results and Discussions

Now let us look at the results based on the  $\tau$ -iteration formula (3.24) and (3.25). Starting from the trial wave functions  $\phi_{ev}$  and  $\phi_+$  defined in (3.9) and (3.6), the energies  $\mathcal{E}_{ev}$  and  $\mathcal{E}_+$  after the first 4 steps of iteration with 5-fold integrations are listed in Tables 1 and 2, together with the energies  $E_{ev,5} = g - \mathcal{E}_{ev,5}$  and  $E_{+,5} = g - \mathcal{E}_{+,5}$ . In Ref.[1] a convergent iteration method to solve the lowest states of the double-well potential, i.e. the  $f$ -iteration, has been given. However, no numerical results have been provided. For comparison the numerical results for  $\mathcal{E}_{ev}$  and  $\mathcal{E}_+$  based on the  $f$ -iteration method is also listed in Tables 1 and 2. For the same folds of integrations the solution of the  $f$ -iteration could only reach a lower accuracy, comparing to the  $\tau$ -iteration, because each step iteration of  $\tau'$  needs only one fold of integration, while each step iteration of  $f$  depends on 2-fold integrations. The accuracy of the obtained energies is higher when  $g$  becomes larger.

Table 1.  $\mathcal{E}_{ev,n}$  and  $E_{ev,5} = g - \mathcal{E}_{ev,5}$

| $g$  | $n$           | 1       | 2       | 3       | 4       | 5       | $E_{ev,5}$ |
|------|---------------|---------|---------|---------|---------|---------|------------|
| 0.05 | $\tau$ -iter. | -0.0341 | -0.0172 | -0.0158 | -0.0163 | -0.0164 | 0.0664     |
| 0.1  | $\tau$ -iter. | -0.0118 | -0.0022 | -0.0016 | -0.0017 | -0.0017 | 0.1017     |
| 0.3  | $\tau$ -iter. | 0.0963  | 0.0973  | 0.0973  | 0.0973  | 0.0973  | 0.2027     |
| 0.5  | $\tau$ -iter. | 0.2035  | 0.2060  | 0.2060  | 0.2060  | 0.2060  | 0.2940     |
| 0.5  | $f$ -iter.    | 0.2036  | 0.2056  | 0.2060  |         |         |            |
| 1    | $\tau$ -iter. | 0.4135  | 0.4310  | 0.4312  | 0.4311  | 0.4311  | 0.5689     |
| 1    | $f$ -iter.    | 0.4135  | 0.4267  | 0.4302  |         |         |            |
| 3    | $\tau$ -iter. | 0.4757  | 0.5105  | 0.5166  | 0.5173  | 0.5173  | 2.4827     |
| 3    | $f$ -iter.    | 0.4757  | 0.5053  | 0.5141  |         |         |            |
| 6    | $\tau$ -iter. | 0.29204 | 0.29399 | 0.29420 | 0.29422 | 0.29422 | 5.70578    |
| 6    | $f$ -iter.    | 0.29204 | 0.29393 | 0.29419 |         |         |            |
| 7    | $\tau$ -iter. | 0.27884 | 0.27957 | 0.27962 | 0.27963 | 0.27963 | 6.72037    |
| 7    | $f$ -iter.    | 0.27884 | 0.27955 | 0.27961 |         |         |            |
| 8    | $\tau$ -iter. | 0.27231 | 0.27265 | 0.27266 | 0.27266 | 0.27266 | 7.72734    |
| 8    | $f$ -iter.    | 0.27231 | 0.27264 | 0.27266 |         |         |            |

Table 2.  $\mathcal{E}_{+,n}$  and  $E_{+,5} = g - \mathcal{E}_{+,5}$ 

| $g$ | $n$           | 1       | 2       | 3       | 4       | 5       | $E_{+,5}$ |
|-----|---------------|---------|---------|---------|---------|---------|-----------|
| 1   | $\tau$ -iter. | 0.4135  | 0.4310  | 0.4312  | 0.4311  | 0.4311  | 0.5689    |
| 1   | $f$ -iter.    | 0.4135  | 0.4267  | 0.4302  |         |         |           |
| 3   | $\tau$ -iter. | 0.3221  | 0.3257  | 0.3258  | 0.3258  | 0.3258  | 2.6742    |
| 3   | $f$ -iter.    | 0.3221  | 0.3254  | 0.3257  |         |         |           |
| 6   | $\tau$ -iter. | 0.27989 | 0.28040 | 0.28041 | 0.28041 | 0.28041 | 5.71959   |
| 6   | $f$ -iter.    | 0.27989 | 0.28039 | 0.28041 |         |         |           |
| 7   | $\tau$ -iter. | 0.27461 | 0.27494 | 0.27494 | 0.27494 | 0.27494 | 6.72506   |
| 7   | $f$ -iter.    | 0.27461 | 0.27493 | 0.27494 |         |         |           |
| 8   | $\tau$ -iter. | 0.27091 | 0.27113 | 0.27113 | 0.27113 | 0.27113 | 7.72887   |
| 8   | $f$ -iter.    | 0.27091 | 0.27113 | 0.27113 |         |         |           |

It is shown clearly that the obtained energies  $E_{ev}$  and  $E_+$  are lower than  $g$  and  $E_{ev} < E_+$ , which are reasonable. When  $g$  increases the two energies become very close to each other and the second step of iteration gives already quite accurate result. It is interesting to notice that for  $g = 1$  the trial function  $\phi_+$  and  $\phi_{ev}$  are the same, therefore starting from  $\phi_+$  the obtained  $\psi_+$  by iteration is the exact ground state wave function for  $g = 1$ .

Table 3.  $\mathcal{E}_{ev}$  and  $E_{ev} = g - \mathcal{E}_{ev}$ 

| $g$  | $\mathcal{E}_{ev}$ | $E_{ev}$ | $g$ | $\mathcal{E}_{ev}$ | $E_{ev}$ |
|------|--------------------|----------|-----|--------------------|----------|
| 0.05 | -0.0164            | 0.0664   | 2.2 | 0.5951             | 1.6049   |
| 0.1  | -0.0017            | 0.1017   | 2.5 | 0.5745             | 1.9255   |
| 0.3  | 0.0973             | 0.2027   | 2.7 | 0.5539             | 2.1461   |
| 0.5  | 0.2060             | 0.2940   | 3.0 | 0.5173             | 2.4827   |
| 0.7  | 0.3065             | 0.3935   | 4.0 | 0.3984             | 3.6016   |
| 1.0  | 0.4311             | 0.5689   | 5.0 | 0.3273             | 4.6727   |
| 1.5  | 0.5598             | 0.9402   | 6.0 | 0.29422            | 5.70578  |
| 1.7  | 0.5851             | 1.1149   | 7.0 | 0.27963            | 6.72037  |
| 2.0  | 0.5990             | 1.4010   | 8.0 | 0.27266            | 7.72734  |

In Table 3 listed are the obtained  $\mathcal{E}_{ev}$  for different  $g$  based on the  $\tau$ -iteration. The obtained  $\mathcal{E}_{ev}$  is negative for very small  $g$  and increases when  $g$  increases from 0.05 to 2.0, then decreases when  $g$  increases further. However, the energy for the ground state  $E_{ev} = g - \mathcal{E}_{ev}$  monotonically increases with increasing  $g$ .

Table 4.  $\mathcal{E}_+$ ,  $E_+ = g - \mathcal{E}_+$  and  $\mathcal{E}_+^N = \sum_{m=0}^N \epsilon_{m+1}/g^m$

| $g$ | $\mathcal{E}_+$ | $E_+$    | $N_{min}$ | $N_{max}$ | $\mathcal{E}_+^N$ | $O(e^{-\frac{4}{3}g})$  |
|-----|-----------------|----------|-----------|-----------|-------------------|-------------------------|
| 1   | 0.4311          | 0.5689   | -         | -         | -                 | $\sim 0.3$              |
| 2   | 0.3664          | 1.6336   | -         | -         | -                 | $\sim 0.07$             |
| 3   | 0.3258          | 2.6742   | -         | -         | -                 | $\sim 0.02$             |
| 4   | 0.3024          | 3.6976   | -         | -         | -                 | $\sim 0.005$            |
| 5   | 0.2888          | 4.7112   | -         | -         | -                 | $\sim 0.001$            |
| 6   | 0.28041         | 5.71959  | 10        | 17        | 0.2807            | $\sim 3 \times 10^{-4}$ |
| 7   | 0.27494         | 6.72506  | 14        | 22        | 0.27501           | $\sim 4 \times 10^{-5}$ |
| 8   | 0.27113         | 7.72887  | 12        | 34        | 0.27115           | $\sim 2 \times 10^{-5}$ |
| 9   | 0.268336        | 8.731664 | 11        | 35        | 0.268339          | $\sim 6 \times 10^{-6}$ |

The energy  $\mathcal{E}_+$  and  $E_+$  are given in Table 4 together with  $\epsilon_+$  calculated from the asymptotic power expansion to  $1/g$ . It can be seen that only when  $g$  is large enough, say  $g \geq 6$ , the asymptotic expansion is meaningful. For such large  $g$  the obtained  $\mathcal{E}_+$  from the iteration and from the power expansion are comparable up to a quite accurate level. The power expansion of  $\mathcal{E}_+$  to  $1/g$  is an asymptotic one. For a fixed and large enough  $g$ -value the summation up to a certain number of terms becomes stable. When increasing the number of summed terms further a plateau of the energy  $\mathcal{E}_+^N$  (within the number of summed terms  $N_{min} < N < N_{max}$  shown in Table 4) is obtained, which gives the  $E_+$ -value accurate to a certain level. However, beyond a certain number of terms ( $N > N_{max}$ ) the result becomes unstable and meaningless. It should be noticed that the asymptotic expansion result within the plateau region does not give the accurate value of the energy. It differs from the iteration one in the order of  $e^{-\frac{4}{3}g}$ , which provides the limitation of the accuracy of the asymptotic expansion.

Recently it has been proved[5] that the  $f$ -iteration in one-dimensional problem is convergent if the trial function is properly chosen to have a finite perturbed potential  $w(x)$ , satisfying the conditions  $w(x) > 0$ ,  $w'(x) < 0$  and  $w(\infty) = 0$ . There is no restriction to the magnitude of  $w(x)$ . For the double-well potential,  $w_{ev}(x)$  defined in (3.10) satisfies these conditions when  $g \geq 1$ . Our numerical results show that both  $f$ - and  $\tau$ -iteration are convergent for  $g \geq 1$ , although it is not an easy task to prove the convergence of the  $\tau$ -iteration. Furthermore, the two iteration series could also be applied in some region of  $g < 1$ , where  $w_{ev}(x)$  is still positive and finite but no more a monotonic function of  $x$ , and reasonable results are obtained. However, for very small  $g$  (e.g.  $g < 0.4$ ), where  $w(x)$  becomes negative in some  $x$ -region, the  $f$ -iteration does not give reasonable results and the obtained  $\mathcal{E}_n$  becomes unstable, while the  $\tau$ -iteration could still work, although the convergence becomes slower when  $g$  is smaller. The reason is the following: The iterative formula for the perturbed wave function  $f$  is expressed as sum of two terms. It could become negative if the term containing  $w(x)$  becomes negative. This would give a negative  $\psi(x)$  in some  $x$ -region, while the solution for the ground state wave function

$\psi(x)$  should be always positive. For the  $\tau$ -iteration the perturbed wave function  $e^{-\tau(x)}$  is always positive for any finite  $\tau(x)$ , either positive or negative. This condition is fulfilled as long as  $|w(x)|$  is finite and  $w(\infty) \rightarrow 0$ . Thus, the  $\tau$ -iteration gives less restrictions to the perturbed potential  $w(x)$ . Therefore, it is of interests to further study the condition for the convergence of the  $\tau$ -iteration and to apply it to other physics problems where perturbation method could not be applied.

## Acknowledgment

The author is grateful to Professor T. D. Lee for his continuous and substantial instructions and advice. This work is partly supported by National Natural Science Foundation of China (NNSFC, No. 20047001).

## Reference

1. R. Friedberg, T. D. Lee, W. Q. Zhao and A. Cimenser, Ann. Phys. 294(2001)67
2. R. Friedberg, T. D. Lee, W. Q. Zhao, Ann. Phys. 288(2001)52
3. Zhao Wei-Qin, Commun. Theoret. Phys. (Beijing,China) 42(2004)37
4. R. Friedberg, T. D. Lee and W. Q. Zhao, IL Nuovo Cimento A112(1999)1195
5. R. Friedberg and T. D. Lee, Ann. Phys. 308(2003)263, quant-ph/0407207

# Appendix

## Appendix A. Asymptotic series expansion of $\tau_+$ and $\mathcal{E}_+$

Starting from the integral equation of  $\tau'_+$  of the first equation in (3.20)

$$\tau'_+(x) = 2\phi_+^{-2}(x) \int_x^\infty \phi_+^2(y) [-(u(y) - \mathcal{E}_+) - \frac{1}{2} (\tau'_+(y))^2] dy \quad (A.1)$$

an asymptotic series expansion of  $\tau_+$  and  $\mathcal{E}_+$  could be obtained. From (A.1) it is easy to obtain the following equation

$$(\frac{1}{2} \phi_+^2 \tau'_+)' = [(u - \mathcal{E}_+) + \frac{1}{2} (\tau'_+)^2] \phi_+^2 \quad (A.2)$$

Using the definition of  $\phi_+$  in (3.6), the above equation leads to

$$gS'_0\tau'_+ = \frac{1}{2} \tau_+'' - S'_1\tau'_+ - (u - \mathcal{E}_+) - \frac{1}{2} (\tau'_+)^2. \quad (A.3)$$

Now let us expand both  $\tau'_+(x)$  and  $\mathcal{E}_+$  in power series of  $g^{-1}$  as following

$$\tau'_+ = \sum_1^\infty \frac{1}{g^m} S'_{m+1} \quad \text{and} \quad \mathcal{E}_+ = \sum_0^\infty \frac{1}{g^m} \epsilon_{m+1}. \quad (A.4)$$

Substituting (A.4) into (A.3), comparing terms proportional to  $g^{-n}$ , we obtain a series of equations for  $\{S_m\}$  and  $\{\epsilon_m\}$ :

$$\begin{aligned} g^0 \quad S'_0 S'_2 &= -(u - \epsilon_1) \\ g^{-1} \quad S'_0 S'_3 &= \frac{1}{2} S_2'' - S'_1 S'_2 + \epsilon_2 \\ g^{-2} \quad S'_0 S'_4 &= \frac{1}{2} S_3'' - S'_1 S'_3 + \epsilon_3 - \frac{1}{2} S_2'^2 \\ &\dots \\ g^{-m} \quad S'_0 S'_{m+2} &= \frac{1}{2} S_{m+1}'' - S'_1 S'_{m+1} + \epsilon_{m+1} - \frac{1}{2} \sum_{n=1}^{m-1} S'_{n+1} S'_{m+1-n}. \end{aligned} \quad (A.5)$$

The above equations are exactly the same as those obtained in Ref.[5], when taking  $\mathcal{E}_+ = -E$  and  $\epsilon_m = -E_m$ . From the first equation of (A.5), considering  $u = \frac{1}{(1+x)^2}$ , we have  $\epsilon_1 = \frac{1}{4}$  when setting  $x = 1$ .

In the following we derive a compact expression of  $\{S'_m\}$  and  $\{\epsilon_m\}$ . Assuming, for  $m \geq 1$ ,

$$S'_{m+1} = \frac{1}{2^{4m}} \xi^2 \sum_{i=0}^{2m-1} \beta_i(m) \xi^i \quad (A.6)$$

where

$$\xi = \frac{2}{1+x}. \quad (\text{A.7})$$

Considering  $S'_1 = \frac{1}{x+1}$ , from the last equation of (A.5) we have

$$\begin{aligned} S'_0 S'_{m+2} &= -\frac{1}{2^{4m+2}} \sum_{l=0}^{2m-1} \beta_l(m) \xi^{l+3} (l+4) \\ &\quad - \frac{1}{2^{4m+2}} \sum_{n=1}^{m-1} \sum_{i=0}^{2n-1} \sum_{j=0}^{2(m-n)-1} 2\beta_i(n) \beta_j(m-n) \xi^{i+j+4} + \epsilon_{m+1}. \end{aligned} \quad (\text{A.8})$$

For the second summation on the right hand side of (A.8), defining  $i+j+1=l$ , we have  $l_{\min}=1$  and  $l_{\max}=2m-1$ , which leads to

$$\begin{aligned} S'_0 S'_{m+2} &= -\frac{1}{2^{4m+2}} \sum_{l=0}^{2m-1} \beta_l(m) \xi^{l+3} (l+4) \\ &\quad - \frac{1}{2^{4m+2}} \sum_{l=0}^{2m-1} \sum_{n=1}^{m-1} \sum_{i=i_{\min}}^{i_{\max}} 2\beta_i(n) \beta_{l-i-1}(m-n) \xi^{l+3} + \epsilon_{m+1}, \end{aligned} \quad (\text{A.9})$$

where

$$\begin{aligned} i_{\min} &= \max(0, l-2(m-n)) \\ i_{\max} &= \min(2n-1, l-1). \end{aligned} \quad (\text{A.10})$$

To fix  $\epsilon_{m+1}$  we put  $x=1$ , then  $\xi=1$  and  $S'_0=1-x^2=0$ . This gives

$$\epsilon_{m+1} = \frac{1}{2^{4m+2}} \sum_{l=0}^{2m-1} \beta_l(m) (l+4) + \frac{1}{2^{4m+2}} \sum_{l=1}^{2m-1} \sum_{n=1}^{m-1} \sum_{i=i_{\min}}^{i_{\max}} 2\beta_i(n) \beta_{l-i-1}(m-n). \quad (\text{A.11})$$

Introducing  $S'_0 = -\frac{4(\xi-1)}{\xi^2}$ ,  $S'_{m+2}$  could be expressed as

$$\begin{aligned} S'_{m+2} &= \frac{1}{2^{4(m+1)}} \xi^2 \frac{1}{\xi-1} \left\{ \sum_{l=0}^{2m-1} \beta_l(m) (l+4) (\xi^{l+3} - 1) \right. \\ &\quad \left. + \sum_{l=1}^{2m-1} \sum_{n=1}^{m-1} \sum_{i=i_{\min}}^{i_{\max}} 2\beta_i(n) \beta_{l-i-1}(m-n) (\xi^{l+3} - 1) \right\}. \end{aligned} \quad (\text{A.12})$$

Applying the equality

$$\xi^{l+3} - 1 = (\xi-1)(\xi^{l+2} + \xi^{l+1} + \dots + 1) = (\xi-1) \sum_{L=0}^{l+2} \xi^L, \quad (\text{A.13})$$

changing the summation order of  $L$  and  $l$  in (A.12) we finally reach the following expression of  $S'_{m+2}$ :



$$\begin{aligned}
S'_{m+2} &= \frac{1}{2^{4(m+1)}} \xi^2 \sum_{L=0}^{2m+1} \xi^L \left\{ \sum_{l=l_1}^{2m-1} \beta_l(m)(l+4) + \sum_{l=l_2}^{2m-1} \sum_{n=1}^{m-1} \sum_{i=i_{min}}^{i_{max}} 2\beta_i(n)\beta_{l-i-1}(m-n) \right\} \\
&= \frac{1}{2^{4(m+1)}} \xi^2 \sum_{L=0}^{2m+1} \beta_L(m+1) \xi^L
\end{aligned} \tag{A.14}$$

with

$$l_1 = \max(0, L-2), \quad l_2 = \max(1, L-2).$$

For  $m=1$ , it is easy to get  $\beta_0(1)=1$  and  $\beta_1(1)=1$ . Finally we obtain, for  $m \geq 1$ ,

$$\beta_L(m+1) = \beta_L^0(m+1) + \Delta\beta_L(m+1) \tag{A.15}$$

where

$$\begin{aligned}
\beta_L^0(m+1) &= \sum_{l=\max(0, L-2)}^{2m-1} \beta_l(m)(l+4) \\
\Delta\beta_L(m+1) &= \sum_{l=\max(1, L-2)}^{2m-1} \sum_{n=1}^{m-1} \sum_{i=i_{min}}^{i_{max}} 2\beta_i(n)\beta_{l-i-1}(m-n)
\end{aligned} \tag{A.16}$$

Taking  $L=0$  in (A.15) and (A.16), from (A.11) we have

$$\epsilon_{m+1} = \frac{1}{2^{4m+2}} \beta_0(m+1). \tag{A.17}$$

Based on (A.15) and (A.16) a revised pyramid structure of  $\beta_l(m)$  could be constructed in a similar way as those in Appendix D of Ref.[1].

$$\begin{array}{cccccc}
\beta_1(1) & \beta_0(1) & & & & m=1 \\
\\
\beta_3^0(2) & \beta_2^0(2) & \beta_1^0(2) & \beta_0^0(2) & & \\
\Delta\beta_3(2) & \Delta\beta_2(2) & \Delta\beta_1(2) & \Delta\beta_0(2) & & \\
\beta_3(2) & \beta_2(2) & \beta_1(2) & \beta_0(2) & & m=2 \\
\\
\beta_5^0(3) & \beta_4^0(3) & \beta_3^0(3) & \beta_2^0(3) & \beta_1^0(3) & \beta_0^0(3) \\
\Delta\beta_5(3) & \Delta\beta_4(3) & \Delta\beta_3(3) & \Delta\beta_2(3) & \Delta\beta_1(3) & \Delta\beta_0(3) \\
\beta_5(3) & \beta_4(3) & \beta_3(3) & \beta_2(3) & \beta_1(3) & \beta_0(3) & m=3 \\
\vdots & & \vdots & & & & 
\end{array} \tag{A.18}$$

By using (A.15) and (A.16), we see that each row  $\beta_l(m+1)$  can be obtained from the row  $\beta_l^0(m+1)$  and  $\Delta\beta_l(m+1)$  while the later two could be obtained from  $\beta_l(m)$  above. For example, using  $\beta_1(1) = \beta_0(1) = 1$ , we have

$$\begin{aligned}
\beta_3^0(2) &= \beta_1(1) \cdot (1+4) = 5, \\
\beta_2^0(2) &= \beta_1(1) \cdot 5 + \beta_0(1) \cdot 4 = 9, \\
\Delta\beta_2(2) &= \Delta\beta_3(2) = 0
\end{aligned}$$

etc. The values of the elements in the pyramid are

$$\begin{array}{cccccccccccl}
\beta_l(1) & & & & 1 & 1 & & & & & m = 1 \\
\beta_l^0(2) & & & 5 & 9 & 9 & 9 & & & & \\
\Delta\beta_l(2) & & & 0 & 0 & 0 & 0 & & & & \\
\beta_l(2) & & & 5 & 9 & 9 & 9 & & & & m = 2 \\
\beta_l^0(3) & 35 & 89 & 134 & 170 & 170 & 170 & & & & \\
\Delta\beta_l(3) & 2 & 6 & 8 & 8 & 8 & 8 & & & & \\
\beta_l(3) & 37 & 95 & 142 & 178 & 178 & 178 & & & & m = 3 \\
\beta_l^0(4) & 333 & 1093 & 2087 & 3155 & 4045 & 4757 & 4757 & 4757 & & \\
\Delta\beta_l(4) & 20 & 76 & 148 & 220 & 256 & 256 & 256 & 256 & & \\
\beta_l(4) & 353 & 1169 & 2235 & 3375 & 4301 & 5013 & 5013 & 5013 & & m = 4 \\
& & \dots & \dots & & & & & & & 
\end{array} \tag{A.19}$$

Correspondingly, the energies are

$$\epsilon_2 = \frac{9}{2^6}, \quad \epsilon_3 = \frac{178}{2^{10}} = \frac{89}{2^9}, \quad \epsilon_4 = \frac{5013}{2^{14}}, \quad \text{etc.} \tag{A.20}$$

## Appendix B

For the convenience of applying the coordinate system  $\{S, \alpha\}$  defined in Eq.(2.10), the definition of some quantities introduced in ref.[2] is given in the following. For the new coordinate system

$$(S, \alpha) = (S, \alpha_1(\mathbf{q}), \alpha_2(\mathbf{q}), \dots, \alpha_{N-1}(\mathbf{q})), \tag{B.1}$$

each point  $\mathbf{q}$  in the  $N$ -dimensional space will now be designated by

$$(S, \alpha_1, \alpha_2, \dots, \alpha_{N-1}), \tag{B.2}$$

instead of  $(q_1, q_2, q_3, \dots, q_N)$ . The corresponding line element can be written as

$$d\vec{\mathbf{q}} = \hat{S} h_S dS + \sum_{j=1}^{N-1} \hat{\alpha}_j h_j d\alpha_j; \tag{B.3}$$

the gradient is given by

$$\nabla = \hat{S} \frac{1}{h_S} \frac{\partial}{\partial S} + \sum_{j=1}^{N-1} \hat{\alpha}_j \frac{1}{h_j} \frac{\partial}{\partial \alpha_j}. \tag{B.4}$$

The kinetic energy operator

$$T = -\frac{1}{2}\nabla^2 \quad (B.5)$$

can be decomposed into two parts:

$$T = T_S + T_\alpha, \quad (B.6)$$

with

$$T_S = -\frac{1}{2h_S h_\alpha} \frac{\partial}{\partial S} \left( \frac{h_\alpha}{h_S} \frac{\partial}{\partial S} \right), \quad (B.7)$$

$$T_\alpha = -\frac{1}{2h_S h_\alpha} \sum_{j=1}^{N-1} \frac{\partial}{\partial \alpha_j} \left( \frac{h_S h_\alpha}{h_j^2} \frac{\partial}{\partial \alpha_j} \right), \quad (B.8)$$

in which

$$h_\alpha = \prod_{j=1}^{N-1} h_j, \quad (B.9)$$

and

$$h_S^2 = [(\nabla S)^2]^{-1}, \quad h_1^2 = [(\nabla \alpha_1)^2]^{-1}, \dots, \quad h_j^2 = [(\nabla \alpha_j)^2]^{-1}, \dots \quad (B.10)$$

The volume element in the  $\mathbf{q}$ -space is now

$$d^N \mathbf{q} = h_S h_\alpha dS d\alpha \quad (B.11)$$

with

$$d\alpha = \prod_{j=1}^{N-1} d\alpha_j. \quad (B.12)$$